



INSTRUCTIONS AFTER BRAZILIAN BUTT LIFT

1. When showering: TAKE OFF ALL THE GAUZE AND LET SOAP (PREFERRABLY ANTI-BACTERIAL) RUN OVER THE INCISIONS. Wear your garment all the time for the first month except when you are in the shower. The second month can just be during the day or at night.
2. Please change any bandages or dressings daily. Only use them if you need them to protect your clothing from drainage. Bandages are optional once the incisions are comfortable.
3. It is normal to experience swelling and bruising in the area of liposuction. Other normal findings are:
 - Areas of numbness on the skin for up to six months
 - A cherry-colored drainage from the incisions during the first few days
 - Redness of the surgical scars for three to six months after surgery (this will slowly fade)
4. Try to stay off your bottom for 6 weeks so the fat can create a new blood supply and stay indefinitely. Use pillows etc to alleviate any pressure directly to the buttocks when sitting or lying down.
5. **Please call Dr. Kaplan at the first sign of:**
 - **Excessive (severe) pain and/or swelling**
 - **Redness, drainage, or odor from the incisions**
 - **Fever or chills**
 - **Shortness of breath**
6. It is extremely important that you do not smoke or have any form of nicotine for a minimum of two weeks after surgery. Smoking will delay the healing process or cause skin loss, which increases your risk of postoperative complications. Also, continue to avoid aspirin and other anti-inflammatory medications (NSAIDS), vitamin E, and herbal medications for one week after surgery.
7. You may resume light activity such as walking and light lifting (up to 5 pounds) within the first few days after surgery. Check with Dr. Kaplan before resuming any strenuous activity such as biking, swimming, aerobics and weightlifting (usually three weeks). Also, when returning to more exercise, start slowly and gradually work up to your daily

routine. Every patient's schedule to returning to normal activity is different. A general rule is that you will naturally limit your level of activity based on the level of discomfort you experience.

8. You may drive after you discontinue your narcotic pain medication and when you are comfortable with your mobility.
9. Usually there are no sutures to be removed. You will have a follow up appointment approximately two weeks after surgery.

Post-Op appointment date	Time